

MATH 130
Final Exam Version A
December 2021

Name: Solutions

UNC Email Address: _____

Instructor: _____

- There are 26 questions on this test:
 - 20 are multiple choice and worth 4 points each;
 - 6 are free response and worth 10 points each.
- Calculators are NOT allowed. Answers are to be given in a form that could be typed into a calculator and use units as necessary. I do NOT need to simplify arithmetic / algebraic expressions.
- I am expected to SHOW ALL WORK on the free response questions; no credit will be given for correct answers without supporting work.
- Partial credit may be awarded on multiple choice questions, but it will be based on my answer only; work may be shown but will not be graded.
- Notation and clarity count. My job is to communicate mathematically and make what I am thinking clear.
- I will sign the Honor Pledge when I am finished or I will let my instructor know of any irregularities with this exam.

I have neither given nor received any unauthorized help on this test
and I have conducted myself within the guidelines of the University
Honor Code.

Pledge: _____

MATH 130 Fall 2021
Final Exam Version A

1. If the point $(2, 3)$ lives on the graph of $y = f(x)$, which point must live on the graph of $y = -2f(x-1) + 2$?

- ☒ $(3, -4)$
☐ $(1, -2)$
☐ $(1, -4)$
☐ $(2, 3)$
☐ $(2, -4)$

$$x-1=2$$

$$x=3$$

$$y = -2 \cdot 3 + 2 = -6 + 2 = -4$$

or: shift $(2, 3)$
 R 1 $\rightarrow (3, 3)$
 reflect & double
 vertical $\rightarrow (3, -6)$
 shift up 2 $\rightarrow (3, -4)$

2. Match the function to the set of equations of its vertical asymptotes. Note, each set of equations can be selected more than once.

C $\sin x$

A. $\{x = \frac{\pi}{2} + \pi k, k \in \mathbb{Z}\}$

C $\cos x$

A $\tan x = \frac{\sin x}{\cos x}$

B. $\{x = \pi k, k \in \mathbb{Z}\}$

B $\csc x = \frac{1}{\sin x}$

A $\sec x = \frac{1}{\cos x}$

C. $\{ \}$ (No asymptotes)


B $\cot x = \frac{\cos x}{\sin x}$

3. Find the area of the following triangle.

- ☐ $\sqrt{3}$
☒ $3\sqrt{3}$
☐ $6\sqrt{3}$
☐ $\frac{\sqrt{3}}{3}$


$$\frac{1}{2} \cdot 2 \cdot 6 \cdot \sin 60^\circ$$

$$= 6 \cdot \frac{\sqrt{3}}{2}$$



4. What is the equation of the hyperbola shown?

- ☐ $\frac{x^2}{4} - y^2 = 1$
☐ $x^2 - \frac{y^2}{4} = 1$
☐ $\frac{x^2}{4} + y^2 = 1$
☐ $\frac{y^2}{4} - x^2 = 1$
☒ $y^2 - \frac{x^2}{4} = 1$
☐ $x^2 + \frac{y^2}{4} = 1$



form: $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$

MATH 130 Fall 2021
Final Exam Version A


5. Is the function $f(x) = 4x^3 \cos(x) \tan(x)$ even, odd, or neither?

- ☐ f is odd because $f(-x) = -f(x)$.
☐ f is even because $f(-x) = -f(x)$.
☐ f is odd because $f(-x) = f(x)$.
☒ f is even because $f(-x) = f(x)$.
☐ f is neither because $f(-x) \neq f(x)$ and $f(-x) \neq -f(x)$.

$$\begin{aligned}
 f(-x) &= 4(-x)^3 \cos(-x) \tan(-x) \\
 &= 4(-x^3)(\cos x)(-\tan x) \\
 &= 4x^3 \cos x \tan x
 \end{aligned}$$

6. Find the exact value of $\cos^{-1}\left(\cos\left(\frac{7\pi}{6}\right)\right)$. $\neq \frac{7\pi}{6}$ as $\frac{7\pi}{6} > \pi$

- ☐ $-\frac{\pi}{6}$
☐ $\frac{\pi}{6}$
☒ $\frac{5\pi}{6}$
☐ $\frac{7\pi}{6}$



7. Which of the following is equivalent to $\frac{1+\cos(2\theta)}{2}$?

- ☐ $\sin^2(\theta)$
☒ $\cos^2(\theta)$
☐ $\tan^2(\theta)$
☐ $2\cos(\theta)\sin(\theta)$


$$\frac{1+2\cos^2\theta-1}{2} = \frac{1+1-2\sin^2\theta}{2} = \frac{1+\cos^2\theta-\sin^2\theta}{2}$$

8. Consider the points with polar coordinates:

A $(-6, \frac{2\pi}{3})$ in Q4 B $(6, \frac{2\pi}{3})$ in Q2 C $(6, -\frac{\pi}{3})$ in Q4 D $(-6, -\frac{\pi}{3})$ in Q2

Which correspond to the point with rectangular (Cartesian) coordinates $(-3, 3\sqrt{3})$?

- ☐ A and C
☐ A and D
☐ B and C
☒ B and D



9. Determine the graph of the curve given by parametric equations $x = 2\sin(3t)$, $y = 1 + \cos(3t)$.

- ☐ A line
☐ A line segment
☐ A parabola
☐ A circle
☒ An ellipse

$$\begin{aligned}
 \frac{x}{2} &= \sin 3t & y-1 &= \cos 3t \\
 \left(\frac{x}{2}\right)^2 + (y-1)^2 &= 1
 \end{aligned}$$

MATH 130 Fall 2021
Final Exam Version A

10. Consider the following function, $f(x) = \begin{cases} x^2 + 5 & \text{if } x < 2 \\ 3x - 2 & \text{if } 2 \leq x < 10 \\ 3 & \text{if } x \geq 10 \end{cases}$.

What are the y -intercept(s) of the graph?

$$f(0) = 3 \cdot 0 - 2$$

- ☒ -2
☐ 3
☐ 5
☐ All of the above.
11. Find the exact value of $\tan^{-1}(-\sqrt{3})$.

$$\frac{y}{x} = \frac{-\sqrt{3}/2}{1/2}$$



- ☒ $-\frac{\pi}{3}$
☐ $-\frac{\pi}{6}$
☐ $\frac{2\pi}{3}$
☐ $\tan^{-1}(-\sqrt{3})$ is undefined.

12. Find the exact value of $\cos(195^\circ)$.

$$\frac{\sqrt{2} - \sqrt{6}}{4}$$

$$195 = 135 + 60$$

$$\cos 135^\circ \cos 60^\circ - \sin 135^\circ \sin 60^\circ$$

$$= -\frac{\sqrt{2}}{2} \cdot \frac{1}{2} - \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{3}}{2}$$

$$= \frac{-\sqrt{2} - \sqrt{6}}{4}$$


$$\frac{\sqrt{2} + \sqrt{6}}{4}$$

$$\frac{\sqrt{6} - \sqrt{2}}{4}$$

$$\frac{-\sqrt{2} - \sqrt{6}}{4}$$

13. Find the equation in polar coordinates of the function graphed below.


- ☐ $r = 2$ centered at origin
☐ $r \sin(\theta) = 2$ $y = 2$
☐ $r \cos(\theta) = 2$ $x = 2$
☐ $r = 2 \sin(\theta)$ center at $(0, 1)$
☒ $r = 2 \cos(\theta)$



MATH 130 Fall 2021
Final Exam Version A

14. A dog runs around part of the circumference of a circle of radius 5 meters. His path makes an angle of 230° with the center of the circle. What distance has he run?

- ☒ $\frac{5 \cdot 23\pi}{18}$ meters
☐ $\frac{25 \cdot 23\pi}{36}$ meters
☐ $25 \cdot 115$ meters
☐ $5 \cdot 230$ meters
☐ $\frac{5}{230}$ meters




$$S = 5m \cdot 230 \cdot \frac{\pi}{180^\circ}$$

$$= \frac{23 \cdot 5}{18} \pi m$$

15. Given a triangle with measurements $A = 40^\circ$, $B = 80^\circ$, and $a = 5$, What is b ?

- ☐ $\frac{3 \sin(60^\circ)}{\sin(40^\circ)}$
☐ $\frac{3 \sin(80^\circ)}{\sin(40^\circ)}$
☐ $\frac{5 \sin(60^\circ)}{\sin(40^\circ)}$
☒ $\frac{5 \sin(80^\circ)}{\sin(40^\circ)}$



$$\frac{b}{\sin 80^\circ} = \frac{5}{\sin 40^\circ}$$

$$b = \frac{5 \cdot \sin 80^\circ}{\sin 40^\circ}$$

16. Find the equation of the directrix for the parabola $-8(x - 2) = (y + 1)^2$.

- ☒ $x = 4$
☐ $x = 0$
☐ $x = -6$
☐ $y = 7$
☐ $y = -1$
☐ $y = -3$



$$4a = 8, a = 2$$

vertex $(2, -1)$; opens left

directrix is $x = 2 + 2 = 4$

17. If $\sin \theta = 0.7$, what is the value of $\sin(\theta + \pi)$?

- ☒ -0.7
☐ -0.3
☐ 0.3
☐ 0.7
☐ $\sqrt{0.91}$

$$= \sin \theta \cos \pi + \cos \theta \sin \pi$$


$$= -\sin \theta + 0$$

$$= -0.7$$

MATH 130 Fall 2021
Final Exam Version A

18. Jamie is measuring the height of a monument on campus. She stands 200 feet away from the monument, and looks up at an angle of 70° to see the top of the monument. How tall is the monument?

- ☐ $200 \sin 70^\circ$ feet
☐ $200 \cos 70^\circ$ feet
☒ $200 \tan 70^\circ$ feet
☐ $200 \cot 70^\circ$ feet



$$\tan 70^\circ = \frac{h}{200'}$$

19. Find and simplify the difference quotient for $f(x) = x^2 - 1$.

- ☐ 1
☒ $2x + h$
☐ $2xh + h^2$
☐ $\frac{2xh + h^2 - 2}{h}$

$$\begin{aligned} \frac{f(x+h) - f(x)}{h} &= \frac{(x+h)^2 - 1 - (x^2 - 1)}{h} \\ &= \frac{x^2 + 2xh + h^2 - 1 - x^2 + 1}{h} = \frac{2xh + h^2}{h} = 2x + h \end{aligned}$$

20. Determine the center of the conic given by $x^2 + 2y^2 - 12x + 8y + 26 = 0$.


- ☒ $(6, -4)$
☐ $(-6, -4)$
☐ $(6, -2)$
☐ $(-6, -2)$

$$\begin{aligned} x^2 - 12x + 6^2 + 2(y^2 + 8y + 4^2) &= 26 + 36 + 32 \\ (x-6)^2 + 2(y+4)^2 &= 94 \\ (6, -4) \end{aligned}$$

21. An airplane flies due north from Columbus to Detroit, a distance of 165 miles, and then turns through an angle of 40° and flies to Toronto, a distance of 200 miles. How far is it directly from Columbus to Toronto? Leave your answer in an exact form that could be typed into a calculator to get an approximation and draw a box around it.

$$x^2 = (165 \text{ mi})^2 + (200 \text{ mi})^2 - 2 \cdot 165 \text{ mi} \cdot 200 \text{ mi} \cdot \cos 140^\circ$$

$$x = \sqrt{165^2 + 200^2 - 400 \cdot 165 \cos 140^\circ} \text{ mi}$$



MATH 130 Fall 2021
Final Exam Version A

22. Given $\sec \theta = \frac{7}{4}$, and $\frac{3\pi}{2} < \theta < 2\pi$, find $\tan \theta$.

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$\tan^2 \theta = \left(\frac{7}{4}\right)^2 - 1$$

$$\tan \theta = \pm \sqrt{\frac{49}{16} - 1} \quad \theta \text{ in } Q4; \tan \theta < 0$$

$$\text{So } \tan \theta = -\sqrt{\frac{33}{16}}$$

$$-\frac{\sqrt{33}}{4}$$

23. Find all solutions to the equation $2 \cos \left(5\theta + \frac{\pi}{3} \right) = -1$.



$$\cos \left(5\theta + \frac{\pi}{3} \right) = -\frac{1}{2}$$

$$5\theta + \frac{\pi}{3} = \frac{2\pi}{3} + 2\pi k \quad \text{or} \quad 5\theta + \frac{\pi}{3} = \frac{4\pi}{3} + 2\pi k$$


$$5\theta = \frac{\pi}{3} + 2\pi k \quad \text{or} \quad 5\theta = \pi + 2\pi k$$

$$\theta = \frac{\pi}{15} + \frac{2\pi}{5}k \quad \text{or} \quad \theta = \frac{\pi}{5} + \frac{2\pi}{5}k$$

$$\frac{\pi}{15} + 2\pi k \quad \text{or} \quad \frac{\pi}{5} + 2\pi k, \text{ for any integer } k$$

MATH 130 Fall 2021
Final Exam Version A

24. The number of hours of daylight over time follows a sinusoidal curve. At location A, the maximum amount of daylight, 15 hours, occurs on the summer solstice which is on day 172. The minimum amount of daylight at location A happens on the winter solstice, when they get 9 hours of daylight on day 355. This repeats annually with a period of 366 days.



$$355 - 366 = -11$$

$$\frac{1}{2} \cdot 366 = 183$$

$$-11 - 183 = -194$$

- (a) If we use the cosine function to model this curve, what is the phase shift in days? (include left or right in your answer)

Right 172 or Left 194

- (b) What is the average amount of daylight for the whole year? (This corresponds to the midline of the function.)

$$\frac{15 + 9}{2} = \frac{24}{2} = 12$$

12 hours

- (c) Write an equation using cosine that gives the hours of daylight y as a function of time t in days. (The solution is graphed above.)

Amp: $\frac{1}{2}(15 - 9) = \frac{1}{2} \cdot 6 = 3$

Period: $366 = \frac{2\pi}{B}$; $B = \frac{2\pi}{366} = \frac{\pi}{183}$

$$y = 3 \cos\left(\frac{\pi}{183}(t - 172)\right) + 12$$

or $y = 3 \cos\left(\frac{\pi}{183}(t + 194)\right) + 12$

MATH 130 Fall 2021
Final Exam Version A

25. Find the inverse function for $f(x) = \frac{3x+2}{4x-1}$ and give the domain and range of f^{-1} in interval notation.

$$x = \frac{3y+2}{4y-1}$$

$$x(4y-1) = 3y+2$$

$$4xy - x = 3y + 2$$

$$4xy - 3y = x + 2$$

$$y(4x-3) = x+2$$

$$y = \frac{x+2}{4x-3} = f^{-1}(x)$$

$$\text{Domain } f^{-1}: x \neq 3/4$$

$$x \neq 1/4 \quad \text{Domain } f = (-\infty, 1/4) \cup (1/4, \infty) \\ = \text{Range } f^{-1}$$

$$f^{-1}(x) = \frac{x+2}{4x-3}$$

$$\text{Domain: } (-\infty, 3/4) \cup (3/4, \infty)$$

$$\text{Range: } (-\infty, 1/4) \cup (1/4, \infty)$$

26. Establish the identity $\tan^2 x = \frac{\sin^2(2x)}{(1 + \cos(2x))^2}$

$$\frac{\sin^2(2x)}{(1 + \cos(2x))^2} = \frac{(2 \sin x \cos x)^2}{(1 + 2 \cos^2 x - 1)^2}$$

$$= \frac{4 \sin^2 x \cos^2 x}{(2 \cos^2 x)^2}$$

$$= \frac{4 \sin^2 x \cos^2 x}{4 \cos^4 x}$$

$$= \frac{\sin^2 x}{\cos^2 x}$$


$$= \tan^2 x$$

MATH 130 Fall 2021
Final Exam Version A

(This page is intentionally left blank to be used for scratch work if necessary.)

MATH 130 Fall 2021
Final Exam Version A

This page will NOT be graded and you do NOT need to fill out the unit circle. It is included in case you find it helpful.



Formulas

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta$$

$$\sin\left(\frac{\theta}{2}\right) = \pm \sqrt{\frac{1 - \cos \theta}{2}}$$

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

$$\text{Area: } \frac{1}{2}ab \sin C$$

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$$

$$= 2 \cos^2 \theta - 1$$

$$= 1 - 2 \sin^2 \theta$$

$$\cos\left(\frac{\theta}{2}\right) = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

Parabola:

$$(y - k)^2 = \pm 4a(x - h)$$

$$(y - k)^2 = 4\rho(x - h)$$

$$(x - h)^2 = \pm 4a(y - k)$$

$$(x - h)^2 = 4\rho(y - k)$$

Ellipse:

$$\frac{(x - h)^2}{a^2} + \frac{(y - k)^2}{b^2} = 1$$

$$\frac{(x - h)^2}{b^2} + \frac{(y - k)^2}{a^2} = 1$$

Hyperbola:

$$\frac{(x - h)^2}{a^2} - \frac{(y - k)^2}{b^2} = 1$$

$$\frac{(y - k)^2}{a^2} - \frac{(x - h)^2}{b^2} = 1$$